



COPING WITH
SUICIDAL THOUGHTS

COUNSELLING AND
PSYCHOLOGICAL
SERVICES (CAPS)



THE UNIVERSITY OF
SYDNEY

If you are thinking about suicide, you are not alone. Many people have thoughts of suicide, for a number of reasons. Thoughts of suicide can be very scary and you may think that nothing can be done to change your situation.

It is important to know that thinking about suicide does not mean that you will lose control or act on these thoughts. Having thoughts of suicide also does not mean that you are weak, or 'crazy'. Many people think about suicide because they are looking for a way to escape the pain they are feeling.

These pages will help you to understand your thoughts and feelings and will provide practical ideas about how to keep safe and how to better manage your suicidal thoughts.

CONTENTS

- 01 Thinking about suicide? What should I do?
- 02 Safety Plan
- 03 How can I better understand my suicidal thoughts and feelings?
- 05 What else can I do to decrease thoughts of suicide?
- 07 How can I decrease chances that I will feel suicidal in the future?
- 08 Useful information

THINKING ABOUT SUICIDE? WHAT SHOULD I DO?

1




If you are thinking about suicide, you are not alone. Many people have thoughts of suicide, for a number of reasons. Thoughts of suicide can be very scary.

You probably feel hurt, confused, overwhelmed and hopeless about your future. You may feel sadness, grief, anger, guilt, shame, or emptiness. You may think that nothing can be done to change your situation. Your feelings may seem like they are just too much to handle right now.

It is important to know that thinking about suicide does not mean that you will lose control or act on these thoughts. Having thoughts of suicide does not mean you are weak, or 'crazy'. Many people think about suicide because they are looking for a way to escape the pain they are feeling.

Even though your situation seems hopeless and you wonder if you can stand another minute of feeling this bad, there are ways to get through this and feel better. You don't have to face this situation alone. Help is available. Here are a few ideas that you can use right now.

- **Connect with others:** If you are worried that you may lose control or do something to hurt yourself, tell someone. Make sure you are around someone you trust. If you live alone, ask a friend or family member to stay with you. If you don't know anyone or can't reach friends or family members call Lifeline: 131-114 or Salvos: 9331-2000.
- **Keep your home safe by getting rid of ways to hurt yourself:** It is important to get rid of things that could be used to hurt or kill yourself, such as pills, razor blades, or guns. If you are unable to do so, go to a place you can feel safe.



- **Develop a safety plan:** It is very helpful to have a written safety plan when you have thoughts of hurting yourself. Have a trusted family member, friend, or professional help you to complete this safety plan. Keep this plan somewhere you can see or find easily. Write down the steps you will take to keep yourself safe (see the following example). Follow the steps. If you follow these steps and still do not feel safe, call a crisis line, get yourself to a hospital emergency department or call 000.

This information is not intended to replace professional care with a therapist/counsellor or physician.

If you have thoughts of hurting yourself, start at Step 1. Go through each step until you are safe. Remember, suicidal thoughts can be very strong. It may seem they last forever. With support and time, these thoughts will usually pass. When they pass, you can put your energy into sorting out problems that have contributed to you feeling so badly. The hopelessness you

may feel now will not last forever. It is important to reach out for help and support. You can get through this difficult time. Since it can be hard to focus and think clearly when you feel suicidal, please print this page and put it in places where you can easily use it, such as your purse, wallet or by the phone.

[Download Safety Plan in PDF format](#)

Do the following activities to calm/comfort myself:
Remind myself of my reasons for living:
Call a friend or family member: Name: _____ Phone: _____
Call a backup person if person above is not available: Name: _____ Phone: _____
Call a health care professional (psychologist, psychiatrist, therapist, local doctor): Name: _____ Phone: _____
Call my local crisis line: Phone: _____
Go somewhere I am safe: Place: _____
Go to the Emergency Department at the nearest hospital.
If I feel that I can't get to the hospital safely, call 000 and request transportation to the hospital. They will send someone to transport me safely.

Some problems and experiences, especially those that have been around a long time, can leave you feeling hopeless and overwhelmed. At these times, you may think that you have no options left. You may think about suicide as a way to escape intense emotional pain.

People who kill themselves often think that their problems are unbearable and can't be fixed. They feel nothing they have tried has or will change their situation. Their emotional pain can distort thinking so it becomes harder to trust, or to see possible solutions to problems, or to connect with available love and support. Even if it seems that you can't stand another minute, it is important to remember that feelings (e.g., grief, anger, sadness, loneliness, shame), especially at this intense level, don't last forever. Sometimes thoughts of suicide can become very strong, especially if you have taken drugs or alcohol. It is important not to use non-prescription drugs or alcohol, particularly when you feel hopeless or are thinking about suicide.

Some of the thoughts you may be having are:

- believing there are no other options
- sensing your family or friends would be better off without you
- thinking you've done something so horrible that suicide is the only option
- experiencing unbearable pain that feels like it will go on forever
- wanting to escape your suffering
- wanting to let your loved ones know how much you hurt, or
- wanting to hurt or get revenge on others

Your feelings of pain are very real. However, it is important to know that there is hope. With



the help of professionals and the support of family and friends, you can learn about what is causing your suffering and how you can change or manage it.

Hurting or killing yourself is not your only option. Professionals can help you learn new skills for dealing with your pain. These might include: developing new skills to cope; seeing your problems in a new light; improving your ability to handle intense and painful emotions; improving your relationships; increasing your social supports; or medications.

Some other things that may lead you to think of suicide are:

- **Mental health problems:** Some mental health problems, such as depression or anxiety, can increase feelings of suicide. Mental health problems are treatable. It is important to talk with your doctor if you feel low, depressed, or anxious. Counselling or medication may help. There are also free resources that can help such as [Mood Gym](#) or workbooks like [Back From the Bluez](#).

- **Conflict with loved ones:** You may feel that your family or friends would be better off without you. It is important to remember that conflict with others doesn't last forever. Ending your life is not a way to solve that conflict. We know that people who lose a loved one to suicide say their lives are not better off.
 - **Loss:** Many different types of loss can increase the chances of feeling suicidal. Some examples that may set off feelings of suicide include: a break-up; losing a job; losing social status; or losing a loved one or friend. Knowing someone who has died by suicide can increase the chance that you think of suicide as an option. As difficult as your loss may seem, there are people and services that can help you get through difficult times, such as Lifeline: 131-114 or Grief Support: 9489-6644.
 - **Financial/legal problems:** Financial or legal problems, such as overwhelming debt, gambling problems, or problems with the law, can be very stressful. It is important to know that there may be free services that can help you deal with financial or legal problems. These include the G-Line (Gambling Support): 1800-633-635.
 - **Lack of connection to friends and others:** Thoughts of suicide can increase if you spend a lot of time alone, or don't feel you can tell anyone your problems. Talk to someone, like a professional, about ways that you can increase social supports in your life. You may feel that the people that are in your life don't understand the pain you are feeling. Talk to a professional about ways that you can let others know of the pain you are feeling.
 - **Drug and alcohol problems:** Using alcohol or drugs can make feelings of depression, anxiety, and thoughts about suicide worse. Drugs and alcohol can change the way you think about problems in your life. If drugs or alcohol are causing your problems you can get information on treatment from the Alcohol and Drug Information (and Referral) Service on 9361-8000. You might also visit justaskus.org.au for online information.
 - **Medical problems:** Medical problems such as diabetes, thyroid problems, pain and other chronic problems can increase chances that you may think about suicide. Make sure you have proper medical care for health problems. Some medications can also increase feelings of suicide. It is important to speak to your doctor about this. The University Health Service bulk bills and can be contacted on 9351-3484 (Wentworth Building) and 9351-4095 (Holme Building).
 - **Sexual identity issues:** People who are lesbian, gay, bisexual, or transgender may have a higher risk of suicide. Confusion about sexual identity and fears of possible or real rejection from family and friends can make things worse. There is support available. The Gay and Lesbian Counselling Service offers phone counselling 7 days from 5:30 to 10:30pm on 8594-9596.
- You can also talk to a counsellor at the University Counselling and Psychological Services about any/all of these types of problems. The service is free, confidential and staffed by registered psychologists. To contact us call 8637 8433 or 8627 8437. **Our website** also has further information, self-help, ideas and links to other useful resources:

WHAT ELSE CAN I DO TO DECREASE THOUGHTS OF SUICIDE?

SOME EXAMPLES OF COMMON PROBLEMS AND IDEAS ARE:

Problem: Depressed mood

Possible solution:

- Call Lifeline on 131-114 for emotional support, short-term problem-solving and referrals for longer term help.
- Speak to a counsellor at the University Counselling and Psychological Services: 8627 8433 or 8627 8437.
- See your family doctor to discuss options for treatment (e.g., medications, changes in medications, undiagnosed illnesses).
- Take care of yourself by resting, exercising regularly, eating regularly and spending time with your friends.

Problem: End of a relationship

Possible solution:

- Talk to friends about the pain you feel.
- Get help from a crisis line or a counsellor at the University Counselling and Psychological Services.
- Join a community social group or a **university club or society**.

Problem-solve: It is always helpful to think of ways other than suicide that you can solve your problems. First, make a list of all the problems you are dealing with in your life. Second, make a list of all the solutions you can think of to those problems. You can ask someone you trust to help you with this. Dealing with one or two small problems can help to put an end to immediate feelings of suicide. Once you are thinking more clearly, you can tackle other bigger problems. For more information you can find an online resource on **problem solving**.

Think of a reason for living: Most people who think about suicide want to escape their pain, but they do not always want to die. When you feel low, it's easy to stay focused on things that are negative and upsetting in your life. This makes it easy to think of suicide as the only option. Start thinking about some reasons you have for living. For example, many people have relationships with loved ones, pets they love, religion, goals and dreams, or responsibilities to others in their life that give them reasons to live and prevent them from acting on their suicidal thoughts. Think of all the reasons you have for living. Write them down. Remind yourself of them when you are feeling low.

Remember things that have helped in the past: Many people have had thoughts of suicide before. Think of some of the things that helped you feel better when you faced the same types of problems in the past. Some examples are: having faith and trust that time always helps; reaching out to friends and family; seeing a professional; going to a support group; following a safety plan; doing something you enjoy; not being alone; keeping a journal; or not drinking or using drugs.



Talk to a trusted friend, family member, or professional: It is important to speak to someone you trust about how you feel. Sometimes just talking about how you feel can help. It is important to be open about all of your thoughts. If you have a suicide plan, it is important to tell someone what your plan is. People often say they are relieved that they shared how they felt with someone. Talking can help you feel less alone.

Get treatment for mental health problems: It is important to get treatment for depression, anxiety, and alcohol and drug problems. Just seeing your family doctor may not be enough. It can help to see a mental health specialist, such as a psychologist or psychiatrist. You can get referrals from your doctor or learn how to find a specialist from one of the referral lines listed on the last page. If you are already receiving treatment, speak up if your treatment plan is not working.

Do the opposite of how you feel: When you have thoughts of suicide, it can be helpful to do the opposite of how you feel. For example, when people feel depressed they usually want to be left alone. Doing the opposite, for example getting in touch with others, can help with feelings of depression.

HOW CAN I DECREASE CHANCES THAT I WILL FEEL SUICIDAL IN THE FUTURE?

Get professional support: Speak to a counsellor at the University Counselling and Psychological Services: 8627 8233 or 8627 8437. Alternatively you can get help and referrals from your doctor or from referral lines listed in the Key Contact Numbers at the top of this page. If the first referral doesn't work for you, ask for another.

Identify high-risk triggers or situations: Think about the situations or factors that increase your feelings of despair and thoughts of suicide. Work to avoid those situations. For example, going to a bar and drinking with friends may increase feelings of depression. If this is a trigger for you, avoid going to a bar or seeing friends who drink.

Self-care: Taking good care of yourself is important to feel better. It is important to do the following:

- eat a healthy diet
- get some exercise every day
- get a good night's sleep, or
- decrease or stop using alcohol or drugs, as these can make feelings of depression and suicide worse.

Follow through with prescribed medications: If you take prescription medications, it is important to make sure you take them as your doctor directed. Speak with your doctor if medications aren't working or if side effects are causing you problems. If you have just begun taking antidepressants, it is important to know that symptoms of depression resolve at different rates. Physical symptoms such as energy or sleep may improve first. Improvement in mood may be delayed. Speak to your doctor if you are feeling worse.

Structure and routine: Keep a regular routine as much as possible, even when your feelings seem out of control. Here are some tips for creating structure in your life:

- wake up at a regular time
- have a regular bed time
- have planned activities in your day, such as going for a walk or going to the gym, or
- continue to go to work or school

Do things you enjoy: When you are feeling very low, do an activity you enjoy. You may find that very few things bring you pleasure. Think of things you used to enjoy doing at times you didn't feel so depressed or suicidal. Do these things, even if they don't bring you enjoyment right now. Giving yourself a break from suicidal thoughts can help, even if it's for a short time.

Think of personal goals: Think of personal goals you have for yourself, or that you've had in the past. Some examples are: to read a particular book; travel; get a pet; move to another place; learn a new hobby; volunteer; go back to school; or start a family.

This document is reproduced and adapted with permission from Dr. Joti Samra and Dr. Dan Bilsker (Lead Authors), Consortium for Organizational Mental Health (COMH; www.comh.ca), CARMHA, Faculty of Health Sciences, Simon Fraser University, Suite 2400, 515 W. Hastings Street, Vancouver, BC V6B 5K3. Inquiries may be directed to info@comh.ca.

KEY CONTACT NUMBERS

Sydney University Counselling and Psychological Services (CAPS)
(02) 86278433 or (02) 8627 8437

University Health Service
(02) 93513484

Mental Health Line (will put you in touch with the Adult Crisis Team in your local area)
1800 011 511

Lifeline (24 Hours)
131114

Salvo Care Line (24 Hours)
(02) 93316000

Domestic Violence Line (24 Hours)
1800200526 or 1800656463

Alcohol and Drug Information (and referral) Service (24 Hours)
(02) 93618000 (Sydney) or
1800422599 (NSW Country)

Sexual Assault (24 hours)
1800424017

G-Line (Gambling)
1800633635

Gay & Lesbian Counselling Service
(02) 85949596

Grief Support (24 Hours)
(02) 94896644

HIV/AIDS Info Line
(02) 93329700

LINKS

[Counselling and Psychological Services \(CAPS\)](#)

[Australian Psychological Society](#)

[Beyond Blue](#)

[Black Dog Institute](#)

[Mood Gym](#)

[Just Ask Us](#)

[Self-help Workbooks](#)

[Other Self-care Workbooks](#)



T +61 2 8627 8433
T +61 2 8627 8437
E caps.admin@sydney.edu.au
sydney.edu.au/counselling



Produced by the University Publishing Service, the University of Sydney, June 2011.
The University reserves the right to make alterations to any information
contained within this publication without notice. UPS51445
© 2011 Counselling and Psychological Services (CAPS), The University of Sydney

ABN 15 211 513 464
CRICOS 00026A