

# 2 Unit Bridging Course - Day 11

## The Logarithm Function

Collin Zheng



# The Logarithm Function $y = \ln x$

Recall from the previous module that if  $y = e^x$ , then

$$x = \ln y.$$

Hence the **Logarithm Function**

$$y = \ln x \quad (\text{for all } x > 0)$$

is precisely the inverse function of the exponential function.

That is,

$$\text{If } f(x) = e^x, \text{ then } f^{-1}(x) = \ln(x)$$

# The Logarithm Function $y = \ln x$

Recall from the previous module that if  $y = e^x$ , then

$$x = \ln y.$$

Hence the **Logarithm Function**

$$y = \ln x \quad (\text{for all } x > 0)$$

is precisely the inverse function of the exponential function.

That is,

$$\text{If } f(x) = e^x, \text{ then } f^{-1}(x) = \ln(x)$$

# The Logarithm Function $y = \ln x$ (cont.)

Recall the *Cancellation Property* for mutually inverse functions  $f$  and  $f^{-1}$ :

$$f^{-1}(f(x)) = x \quad \text{and} \quad f(f^{-1}(x)) = x.$$

Setting  $f(x) = e^x$  and  $f^{-1}(x) = \ln(x)$ , we thus have the following two identities:

$$\ln(e^x) = x$$

$$e^{\ln x} = x \quad \text{for all } x > 0$$

That is, 'exponentiating' and 'logging' are mutually inverse operations and therefore cancel each other out.

## The Logarithm Function $y = \ln x$ (cont.)

Recall the *Cancellation Property* for mutually inverse functions  $f$  and  $f^{-1}$ :

$$f^{-1}(f(x)) = x \quad \text{and} \quad f(f^{-1}(x)) = x.$$

Setting  $f(x) = e^x$  and  $f^{-1}(x) = \ln(x)$ , we thus have the following two identities:

$$\ln(e^x) = x$$

$$e^{\ln x} = x \quad \text{for all } x > 0$$

That is, 'exponentiating' and 'logging' are mutually inverse operations and therefore cancel each other out.

## Example

We can use this cancelling phenomenon to simplify  $e^{2\ln x}$ :

$$\begin{aligned} e^{2\ln x} &= e^{\ln x^2} && \longleftarrow \text{using log law \#3} \\ &= x^2. && \longleftarrow \text{using cancellation property} \end{aligned}$$

## Example

We can use this cancelling phenomenon to simplify  $e^{2\ln x}$ :

$$\begin{aligned} e^{2\ln x} &= e^{\ln x^2} && \longleftarrow \text{using log law \#3} \\ &= x^2. && \longleftarrow \text{using cancellation property} \end{aligned}$$

## Example

We can use this cancelling phenomenon to simplify  $e^{2\ln x}$ :

$$\begin{aligned} e^{2\ln x} &= e^{\ln x^2} && \longleftarrow \text{using log law \#3} \\ &= x^2. && \longleftarrow \text{using cancellation property} \end{aligned}$$

We can also exploit the cancelling phenomenon to help solve equations.

## Example

For instance, suppose we want to solve  $e^{2x-3} = 7$ .

Logging both sides, we get

$$2x - 3 = \ln 7.$$

Isolating  $x$ , we hence have

$$x = \frac{\ln 7 + 3}{2} \approx 2.47.$$

We can also exploit the cancelling phenomenon to help solve equations.

## Example

For instance, suppose we want to solve  $e^{2x-3} = 7$ .

Logging both sides, we get

$$2x - 3 = \ln 7.$$

Isolating  $x$ , we hence have

$$x = \frac{\ln 7 + 3}{2} \approx 2.47.$$

We can also exploit the cancelling phenomenon to help solve equations.

## Example

For instance, suppose we want to solve  $e^{2x-3} = 7$ .

Logging both sides, we get

$$2x - 3 = \ln 7.$$

Isolating  $x$ , we hence have

$$x = \frac{\ln 7 + 3}{2} \approx 2.47.$$

## Example

A final example – recall our population model from Day 8. That is, the population,  $P(t)$ , of an outback town is growing exponentially according to the formula

$$P(t) = 1000 e^{0.2t},$$

where  $t$  is the number of years after the year 2000.

Recall that the model gave a population of 1000 in the year 2000, 7389 in 2010, and an estimate of roughly 55000 in 2020.

When will the population reach 1,000,000 people?

## Example

A final example – recall our population model from Day 8. That is, the population,  $P(t)$ , of an outback town is growing exponentially according to the formula

$$P(t) = 1000 e^{0.2t},$$

where  $t$  is the number of years after the year 2000.

Recall that the model gave a population of 1000 in the year 2000, 7389 in 2010, and an estimate of roughly 55000 in 2020.

When will the population reach 1,000,000 people?

## Example

A final example – recall our population model from Day 8. That is, the population,  $P(t)$ , of an outback town is growing exponentially according to the formula

$$P(t) = 1000 e^{0.2t},$$

where  $t$  is the number of years after the year 2000.

Recall that the model gave a population of 1000 in the year 2000, 7389 in 2010, and an estimate of roughly 55000 in 2020.

**When will the population reach 1,000,000 people?**

## The Logarithm Function $y = \ln x$ (cont.)

We must set  $P(t)$  to be 1000000 and solve for  $t$  in the equation

$$1000000 = 1000 e^{0.2t}, \text{ i.e. } 1000 = e^{0.2t}.$$

Logging both sides, we get

$$\ln 1000 = 0.2t,$$

so

$$t = \frac{\ln 1000}{0.2} \approx 34.5.$$

That is, the population should reach 1 million by the middle of the year 2034.

## The Logarithm Function $y = \ln x$ (cont.)

We must set  $P(t)$  to be 1000000 and solve for  $t$  in the equation

$$1000000 = 1000 e^{0.2t}, \text{ i.e. } 1000 = e^{0.2t}.$$

Logging both sides, we get

$$\ln 1000 = 0.2t,$$

so

$$t = \frac{\ln 1000}{0.2} \approx 34.5.$$

That is, the population should reach 1 million by the middle of the year 2034.

## The Logarithm Function $y = \ln x$ (cont.)

We must set  $P(t)$  to be 1000000 and solve for  $t$  in the equation

$$1000000 = 1000 e^{0.2t}, \text{ i.e. } 1000 = e^{0.2t}.$$

Logging both sides, we get

$$\ln 1000 = 0.2t,$$

so

$$t = \frac{\ln 1000}{0.2} \approx 34.5.$$

That is, the population should reach 1 million by the middle of the year 2034.

## Practice Questions

- ▶ Simplify  $e^{\frac{1}{2} \ln(x+9)}$ .
- ▶ Simplify  $\ln(e^{2x+1})$ .
- ▶ Solve  $e^{-\ln x} = 2$ .
- ▶ Solve  $\ln 3x = 2$ .
- ▶ Solve  $\ln x^4 - \ln x = 0$ .

## Answers

▶  $\sqrt{x + 9}$ .

▶  $2x + 1$ .


▶  $\frac{1}{2}$ .

▶  $\frac{e^2}{3}$ .

▶ 1.

# The graph of $y = \ln x$

Finally, recall that the graphs of two mutually inverse functions  $f$  and  $f^{-1}$  are symmetric about the diagonal line  $y = x$ :



- ▶  $y = e^x$  and  $y = \ln x$  are mutually inverse functions.  
Hence  $\ln(e^x) = x$  and  $e^{\ln x} = x$ .
- ▶ The graph of  $y = \ln x$  is the reflection of the graph of  $y = e^x$  about the diagonal  $y = x$ .